

SOLAS Session 10: Degenerative

A Case Control Study Of Minimal Invasive Transforaminal Interbody Fusion (MI-TLIF) Versus Extreme Lateral Interbody Fusion (ELIF) At The L4/5 Level – Clinical Outcome And Complications

K. Lam, C. De La Torre; Orthopedics, London Bridge Hospital, London/United Kingdom

Can Anterior Lumbar Surgery Be Performed Safely In An Outpatient Setting?

S. Stephan, C. Johnson, A. Wong, K. Siddique, B. Perri, P. Hill, R. Rao, W. Wagner, N. Anand, B. Khanderoo, E. Nomoto; Orthopaedic Surgery, Cedars Sinai Medical Center, Los Angeles, CA/United States of America

Slow Progression Of Bony Fusion Of Allograft In Transpsoas Lateral Lumbar Interbody Fusion

K. Satake¹, T. Kanemura^{1,2}, K. Ito^{1,2}, S. Tanaka^{1,2}, H. Nakashima³, J. Ouchida³; ¹Spine Center, Konan Kosei Hospital, Konan/Japan, ²Konan Kosei Hospital, Konan/Japan, ³Department of Orthopaedics, Nagoya University Graduate school of Medicine, Nagoya/Japan

Extreme Lateral Interbody Fusion (XLIF) At L5-S1: Feasibility, Indications, And Preliminary Results In A Series Of 31 Cases

V. Stifano¹, R. Roperto², F. Calvosa², F. Boccacci¹, L. Mastronardi²; ¹Neurosurgery, Catholic University of Rome, Rome/Italy, ²Neurosurgery, San Filippo Neri Hospital, Rome/Italy

Comparison Of Foraminal Indirect Decompression Outcomes In Degenerative Spondylolisthesis And Other Degenerative Lumbar Disease After Minimally Invasive Extreme Lateral Interbody Fusion

C. De La Torre¹, K. Lam²; ¹Orthopedics, London Bridge Hospital, London/United Kingdom, ²London Bridge Hospital, London/United Kingdom

Clinical Outcomes After Minimally Invasive Extreme Lateral Interbody Fusion In Degenerative Spondylolisthesis

C. De La Torre, K. Lam; Orthopedics, London Bridge Hospital, London/United Kingdom

Comparison Of Clinical Outcomes And Complications After Minimally Invasive Extreme Lateral Interbody Fusion Based On Body Mass Index

C. De La Torre, K. Lam, K. Uotani, H. Ouyang; Orthopedics, London Bridge Hospital, London/United Kingdom

Fusion Rate For Stand-Alone Lateral Lumbar Interbody Fusion: A Systematic Review

M. Mazur¹, M. Steinhaus², S. Virk², B. Jivanelli², A. Vaishnav², S. McAnany², S. Iyer², T. Albert², C.H. Gang², S. Qureshi²; ¹Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, PA/United States of America, ²Hospital for Special Surgery, New York, NY/United States of America

A Comparison Of Lateral Lumbar Interbody Fusion With Percutaneous Pedicle Screw Fixation Versus Direct Decompression With Posterior Lumbar Fusion: A Propensity-Matched Analysis

J. Ouchida¹, T. Kanemura², K. Satake², K. Ito², S. Tanaka², S. Imagama¹; ¹Department of Orthopaedics, Nagoya University Graduate school of Medicine, Nagoya/Japan, ²Spine Center, Konan Kosei Hospital, Konan/Japan

Durability Of Stand-Alone Lateral Lumbar Interbody Fusion: 5 Year Outcomes

N. Agarwal, M. White, A. Ozpinar, N. Alan, D. Okonkwo, D. Hamilton, A. Kanter; Neurological Surgery, University of Pittsburgh Medical Center, Pittsburgh, PA/United States of America

Radiographic Predictors Of Indirect Neural Decompression In Minimally Invasive Transpoas Lateral Interbody Fusion

C. Walker¹, D. Xu², J. Godzik³, J. Turner³, J. Uribe¹; ¹Barrow Neurological Institute, Phoenix, AZ/United States of America, ²Neurosurgery, Baylor, Houston, TX/United States of America, ³Barrow Brain and Spine, Phoenix, AZ/United States of America